arXiv:hep-ph/0107163v3 31 Jul 2001

DPNU-01-13

$B \to \phi K$ decays in perturbative QCD approach

S. Mishima*
Department of Physics, Nagoya University, Nagoya 464-8602, Japan

We calculate the branching ratios and CP asymmetries of the $B \to \phi K$ decays using perturbative QCD approach, which includes k_T and threshold resummations. Our results of branching ratios are consistent with the experimental data and larger than those obtained from the naive factorization assumption and the QCD-improved factorization approach.

I. INTRODUCTION

Recently the branching ratios of the $B \to \phi K$ decays have been measured by the BaBar [1], BELLE [2] and CLEO [3] collaborations. These decay modes are important to understand penguin dynamics, because the $B \to \phi K$ modes are pure penguin processes.

There is a following problem related a penguin contribution for amplitude [4]. A naive estimate of the loop diagram leads to $P/T \sim \alpha_s/(12\pi)\log(m_t^2/m_c^2) \sim O(0.01)$ where P is a penguin amplitude and T is a tree amplitude. But experimental data for $\mathrm{Br}(B \to K\pi)$ and $\mathrm{Br}(B \to \pi\pi)$ leads to $P/T \sim O(0.1)$. The penguin contribution is enhanced dynamically. This enhancement can not be explained by factorization assumption for nonleptonic two-body B meson decays [5]. This problem is explained by Keum, Li and Sanda using perturbative QCD (PQCD) approach [6].

PQCD method for inclusive decays was developed by many researcher over many years, and this formalism has been successful. Recently, PQCD has been applied to the exclusive B meson decays, $B \to K\pi$ [6], $\pi\pi$ [7], $\pi\rho$, $\pi\omega$ [8], KK [9] and $K\eta^{(\prime)}$ [10]. We calculate the branching ratios for the $B \to \phi K$ modes using this approach to leading order in α_s and leading power in $1/M_B$ where M_B is the B meson mass. The $B \to \phi K$ modes depend only on penguin amplitudes, and the branching ratio for this modes is proportional to $|P|^2$. Thus the $B \to \phi K$ modes are useful to seeing the size of the penguin amplitudes.

In this paper we compute the branching ratio for the $B \to \phi K$ modes using PQCD approach. In Sec. II, we introduce the PQCD formalism for exclusive B meson decays. In Sec. III, we calculate analytic formula of the factorizable amplitudes and the nonfactorizable amplitudes of the decays. In Sec. IV, we predict the branching ratios of the decays and the direct CP asymmetry of the charged mode. Our predicted branching ratios agree with the current experimental data and are larger than the values obtained from the naive factorization assumption and the QCD-improved factorization [11] [12]. In the Appendix, we review the meson wave functions with up to twist-3 terms which are proportional to m_{0K}/M_B or M_ϕ/M_B.

---

*e-mail: mishima@eken.phys.nagoya-u.ac.jp

II. PQCD FACTORIZATION THEOREM

The effective Hamiltonian for $\Delta S = 1$ transitions is

$$H_{\rm eff} = \frac{G_F}{\sqrt{2}} \sum_{q=u,c} V_{qs}^* V_{qb} \left[C_1(\mu) O_1^{(q)}(\mu) + C_2(\mu) O_2^{(q)}(\mu) + \sum_{i=3}^{10} C_i(\mu) O_i(\mu) \right] , \tag{1}$$

where V_{qs}^* and V_{qb} are the Cabibbo-Kobayashi-Maskawa matrix elements [13] and O_{1-10} are local four-fermi operators. The local operators are given by

$$\begin{aligned}
O_1^{(q)} &= (\bar{s}_i q_j)_{V-A} (\bar{q}_j b_i)_{V-A} , & O_2^{(q)} &= (\bar{s}_i q_i)_{V-A} (\bar{q}_j b_j)_{V-A} , \\
O_3 &= (\bar{s}_i b_i)_{V-A} \sum_q (\bar{q}_j q_j)_{V-A} , & O_4 &= (\bar{s}_i b_j)_{V-A} \sum_q (\bar{q}_j q_i)_{V-A} , \\
O_5 &= (\bar{s}_i b_i)_{V-A} \sum_q (\bar{q}_j q_j)_{V+A} , & O_6 &= (\bar{s}_i b_j)_{V-A} \sum_q (\bar{q}_j q_i)_{V+A} , \\
O_7 &= \frac{3}{2} (\bar{s}_i b_i)_{V-A} \sum_q e_q (\bar{q}_j q_j)_{V+A} , & O_8 &= \frac{3}{2} (\bar{s}_i b_j)_{V-A} \sum_q e_q (\bar{q}_j q_i)_{V+A} , \\
O_9 &= \frac{3}{2} (\bar{s}_i b_i)_{V-A} \sum_q e_q (\bar{q}_j q_j)_{V-A} , & O_{10} &= \frac{3}{2} (\bar{s}_i b_j)_{V-A} \sum_q e_q (\bar{q}_j q_i)_{V-A} ,
\end{aligned} \tag{2}$$

where $i,\ j$ is the color indices and q is taken $u,\ d,\ s$ and c. The Wilson coefficients C_{1-10} are calculated up to next-to-leading order by Buras *et al.* [14].

PQCD approach is based on the three scale factorization theorem [15] [16] in which the matrix element can be written as the convolution of the hard part H with the hard scale $t \sim O(M_B)$, the meson wave function Φ with the soft scale $1/b \sim \Lambda_{\rm QCD}$ and the Wilson coefficient C whose evolves from the W boson mass M_W down to the scale t. b is the conjugate variable of the parton transverse momenta k_T. For instance, the $B \to K$ transition form factor is written as

$$F^{BK} \sim \int [dx][db] C(t) \Phi_K(x_2, b_2) H(t) \Phi_B(x_1, b_1) \exp\left[- \sum_{i=1,2} \int_{1/b_i}^{t} \frac{d\bar{\mu}}{\bar{\mu}} \gamma_\phi(\alpha_s(\bar{\mu})) \right] , \tag{3}$$

where x is the momentum fraction of parton and γ_ϕ is the anomalous dimension of meson. The hard part H can be calculated perturbatively and the Feynman diagrams are described as Fig. 1 in leading order of α_s.



FIG. 1. The $B \to K$ transition form factor in leading order of α_s

It is important that the scale of the Wilson coefficient is determined dynamically, since the scale depends on x and b. Then in PQCD, the scale of the Wilson coefficient can reach lower scale than $M_B/2$ which are usually taken in the naive factorization assumption case. The Wilson coefficients for the penguin operator are enhanced as the scale evolves to lower scale. Therefore, the penguin amplitudes are larger than those in the naive factorization assumption.

In this method, we can calculate not only factorizable amplitudes but also nonfactorizable and annihilation amplitudes which can not be calculated in the factorization assumption.

This $B \to K$ transition form factor has two types of double logarithmic corrections in infrared divergences. The one is $\alpha_s \ln^2(Pb)$ from the overlap of collinear and soft divergences which are generated by the correction of the meson wave function as Fig. 2(a) and Fig. 2(b). The resummation of this double logarithms leads to the factor $\exp[-s(P,b)]$ [17] [18]. The explicit form of this factor is shown, for example, in Ref. [19]. This resummation is called k_T resummation. The other double logarithm is $\alpha_s \ln^2(1/x)$ from the end-point region of the momentum fraction x [20]. This double logarithm is generated by the corrections of the hard part as Fig. 2(c) and Fig. 2(d). The resummation of this double logarithms is called the threshold resummation. We use the approximate form of this resummation factor that is $T(x) = N_t\{x(1-x)\}^c$ where $N_t = 1.775$ and $c = 0.3$ [21]. The Sudakov factor from k_T and threshold resummations suppress large b region and end-point of x where the hard part H is singular, the hard part H can then be calculated perturbatively.



FIG. 2. These are the diagrams which generate double logarithms corrections in infrared divergences. Fig. 2(a) and Fig. 2(b) are the correction of the meson wave function. Fig. 2(c) and Fig. 2(c) are the correction of the hard part.

In PQCD approach, we can calculate hard part perturbatively, and we use the meson wave functions which are formed in the light-cone QCD sum rules. The hard part depends on processes, but the wave functions are independent of those. We can determine the parameters in the wave functions, since there are many modes in the B meson two-body decays. So, we can predict quantitative values for the other decays.

In this study, we calculate the hard part in leading order of α_s. The meson wave functions are used with leading twist and twist-3 terms which are proportional to m_{0K}/M_B or M_ϕ/M_B. We review the meson wave functions in the Appendix.

III. $B \to \phi K$ AMPLITUDES

In the light-cone coordinate, the B meson momentum P_1, the K meson momentum P_2 and the ϕ meson momentum P_3 are taken to be

$$P_1 = \frac{M_B}{\sqrt{2}}(1,1,\mathbf{0}_T)\ ,\quad P_2 = \frac{M_B}{\sqrt{2}}(1-r_\phi^2,0,\mathbf{0}_T)\ ,\quad P_3 = \frac{M_B}{\sqrt{2}}(r_\phi^2,1,\mathbf{0}_T)\ , \tag{4}$$

where $r_\phi = M_\phi/M_B$, and the K meson mass is neglected. In this situation, the B meson is at rest. The quark and anti-quark momenta in the mesons are displayed in Fig. 3. Their components are

$$\begin{aligned}
k_1 &= (0, x_1 P_1^-, \mathbf{k}_{1T}), \qquad P_1 - k_1 = (P_1^+, (1-x_1)P_1^-, -\mathbf{k}_{1T})\ ,\\
k_2 &= (x_2 P_2^+, 0, \mathbf{k}_{2T}), \qquad P_2 - k_2 = ((1-x_2)P_2^+, 0, -\mathbf{k}_{2T})\ ,\\
k_3 &= (0, x_3 P_3^-, \mathbf{k}_{3T}), \qquad P_3 - k_3 = (0, (1-x_3)P_3^-, -\mathbf{k}_{3T})\ .
\end{aligned} \tag{5}$$

We can integrate over k_1^+ and choose $k_1^+ = 0$, since the hard part is independent of k_1^+.



(a) (b)

FIG. 3. Momentum assignment

The ϕ meson longitudinal polarization vector ϵ_ϕ and two transverse polarization vector $\epsilon_{\phi T}$ are given by

$$\epsilon_\phi = \frac{1}{\sqrt{2}r_\phi}(-r_\phi^2, 1, \mathbf{0}_T) , \qquad \epsilon_{\phi T} = (0, 0, \mathbf{1}_T) . \tag{6}$$

The B meson wave function for incoming state and the K and ϕ meson wave functions for outgoing state with up to twist-3 terms are written as

$$\Phi^{(in)}_{B,\alpha\beta,ij} = \frac{i\delta_{ij}}{\sqrt{2N_c}} \int dx_1 d^2\mathbf{k}_{1T} e^{-i(x_1 P_1^- z_1^+ - \mathbf{k}_{1T}\mathbf{z_1}_T)} \left[(\not P_1 + M_B)\gamma_5 \phi_B(x_1, \mathbf{k}_{1T})\right]_{\alpha\beta} , \tag{7}$$

$$\Phi^{(out)}_{K,\alpha\beta,ij} = \frac{-i\delta_{ij}}{\sqrt{2N_c}} \int_0^1 dx_2 e^{ix_2 P_2\cdot z_2} \gamma_5 \left[\not P_2 \phi_K^A(x_2) + m_{0K}\phi_K^P(x_2) + m_{0K}(\not v \not n - 1)\phi_K^T(x_2)\right]_{\alpha\beta} , \tag{8}$$

$$\Phi^{(out)}_{\phi,\alpha\beta,ij} = \frac{\delta_{ij}}{\sqrt{2N_c}} \int_0^1 dx_3 e^{ix_3 P_3\cdot z_3} \left[M_\phi \not\epsilon_\phi \phi_\phi(x_3) + \not\epsilon_\phi \not P_3 \phi_\phi^t(x_3) + M_\phi \phi_\phi^s(x_3)\right]_{\alpha\beta} , \tag{9}$$

where i and j is the color indices, α and β are the Dirac indices. m_{0K} is related to chiral symmetry breaking scale, $m_{0K} = M_K^2/(m_d + m_s)$. v and n are defined as $v^\mu = P_2^\mu/P_2^+$ and $n^\mu = z_2^\mu/z_2^- = (0, 1, \mathbf{0}_T)$. We neglect the wave functions which are proportional to the transverse polarization vector ϵ_ϕ^T, because these terms do not appear in our calculations. It must be noted that the expression of $\Phi_\phi^{(out)}$ depends on how to define the sign of ϵ_ϕ. The explicit form of these wave functions will be shown in Sec. IV and the detail is shown in Appendix.

The decay rates for the $B \to \phi K$ have the expressions

$$\Gamma = \frac{G_F^2}{32\pi M_B}|\mathcal{A}|^2 . \tag{10}$$

The decay amplitudes $\mathcal{A}$ and $\bar{\mathcal{A}}$ corresponding to $B^0 \to \phi K^0$ and $\bar{B}^0 \to \phi \bar{K}^0$ respectively, are written as

$$\mathcal{A} = f_\phi V_{ts}V_{tb}^* F_e^P + V_{ts}V_{tb}^* \mathcal{M}_e^P + f_B V_{ts}V_{tb}^* F_a^P + V_{ts}V_{tb}^* \mathcal{M}_a^P , \tag{11}$$

$$\bar{\mathcal{A}} = f_\phi V_{ts}^* V_{tb} F_e^P + V_{ts}^* V_{tb} \mathcal{M}_e^P + f_B V_{ts}^* V_{tb} F_a^P + V_{ts}^* V_{tb} \mathcal{M}_a^P . \tag{12}$$

F_e is the amplitude for the factorizable diagrams which are considered in the factorization assumption. F_a and $\mathcal{M}$ are the annihilation factorizable and the nonfactorizable diagrams which are neglected in the factorization assumption. The indices e and a denote the tree topology as Fig. 3(a) and annihilation topology as Fig. 3(b) respectively. The index P denotes the contribution from the diagram with a penguin operator. The decay amplitudes $\mathcal{A}^+$ and $\mathcal{A}^-$ corresponding to $B^+ \to \phi K^+$ and $B^- \to \phi K^-$ respectively, are written as

$$\mathcal{A}^+ = f_\phi V_{ts}V_{tb}^* F_e^P + V_{ts}V_{tb}^* \mathcal{M}_e^P + f_B V_{ts}V_{tb}^* F_a^P + V_{ts}V_{tb}^* \mathcal{M}_a^P - f_B V_{us}V_{ub}^* F_a^T - V_{us}V_{ub}^* \mathcal{M}_a^T , \tag{13}$$

$$\mathcal{A}^- = f_\phi V_{ts}^* V_{tb} F_e^P + V_{ts}^* V_{tb} \mathcal{M}_e^P + f_B V_{ts}^* V_{tb} F_a^P + V_{ts}^* V_{tb} \mathcal{M}_a^P - f_B V_{us}^* V_{ub} F_a^T - V_{us}^* V_{ub} \mathcal{M}_a^T , \tag{14}$$

where there are very small contribution from tree diagrams, the index T denotes these contribution.

A. Factorizable Amplitudes



FIG. 4. Feynman diagrams contributing to factorizable amplitudes

The factorizable amplitudes which come from Fig. 4 are written as

$$\begin{aligned}
F_e^P &= F_{3e}^P + F_{4e}^P + F_{5e}^P \\
&= 8\pi C_F M_B^4 \int_0^1 dx_1 dx_2 \int_0^\infty b_1 db_1 b_2 db_2 \phi_B(x_1, b_1) \\
&\quad \times \Big\{ \left[(1+x_2)\phi_K^A(x_2) + r_K(1-2x_2)\left(\phi_K^P(x_2) + \phi_K^T(x_2)\right)\right] \\
&\qquad \times E_e(t_e^{(1)}) N_t\{x_2(1-x_2)\}^c h_e(x_1, x_2, b_1, b_2) \\
&\qquad + 2r_K \phi_K^P(x_2) E_e(t_e^{(2)}) N_t\{x_1(1-x_1)\}^c h_e(x_2, x_1, b_2, b_1) \Big\} ,
\end{aligned} \tag{15}$$

$$\begin{aligned}
F_a^P &= F_{a4}^P + F_{a6}^P , \\
F_{a4}^P &= 8\pi C_F M_B^4 \int_0^1 dx_2 dx_3 \int_0^\infty b_2 db_2 b_3 db_3 \\
&\quad \times \Big\{ \left[x_3 \phi_K^A(x_2)\phi_\phi(x_3) - 2r_K r_\phi \phi_K^P(x_2)\left(\left(\phi_\phi^t(x_3) - \phi_\phi^s(x_3)\right) - x_3\left(\phi_\phi^t(x_3) + \phi_\phi^s(x_3)\right)\right)\right] \\
&\qquad \times E_{a4}(t_a^{(1)}) N_t\{x_3(1-x_3)\}^c h_a(1-x_2, x_3, b_2, b_3) \\
&\qquad - \left[(1-x_2)\phi_K^A(x_2)\phi_\phi(x_3) + 2r_K r_\phi \left(2\phi_K^P(x_2) - x_2\left(\phi_K^P(x_2) - \phi_K^T(x_2)\right)\right)\phi_\phi^s(x_3)\right] \\
&\qquad \times E_{a4}(t_a^{(2)}) N_t\{x_2(1-x_2)\}^c h_a(x_3, 1-x_2, b_3, b_2) \Big\} ,
\end{aligned} \tag{16}$$

$$\begin{aligned}
F_{a6}^P &= 16\pi C_F M_B^4 \int_0^1 dx_2 dx_3 \int_0^\infty b_2 db_2 b_3 db_3 \\
&\quad \times \Big\{ \left[2r_K \phi_K^P(x_2)\phi_\phi(x_3) - r_\phi x_3 \phi_K^A(x_2)\left(\phi_\phi^t(x_3) - \phi_\phi^s(x_3)\right)\right] \\
&\qquad \times E_{a6}(t_a^{(1)}) N_t\{x_3(1-x_3)\}^c h_a(1-x_2, x_3, b_2, b_3) \\
&\qquad + \left[r_K(1-x_2)\left(\phi_K^P(x_2) + \phi_K^T(x_2)\right)\phi_\phi(x_3) + 2r_\phi \phi_K^A(x_2)\phi_\phi^s(x_3)\right] \\
&\qquad \times E_{a6}(t_a^{(2)}) N_t\{x_2(1-x_2)\}^c h_a(x_3, 1-x_2, b_3, b_2) \Big\} .
\end{aligned} \tag{17}$$

The tree amplitude F_a^T is the same as F_{a4}^P, but the Wilson coefficient is different. $N_t\{x(1-x)\}^c$ is the factor for the threshold resummation. N_t is a normalization factor and c is a constant, we use $N_t = 1.775$ and $c = 0.3$. The evolution factors are defined by

$$E_e(t) = \alpha_s(t) a_e(t) \exp[-S_B(t) - S_K(t)] \ , \quad E_{ai}(t) = \alpha_s(t) a_{ai}(t) \exp[-S_K(t) - S_\phi(t)] \ . \tag{18}$$

The explicit forms of the factor $S_i(t)$ are given, for example, in Ref. [6]. The hard scales t, which are the typical scales in hard process, are given by

$$\begin{aligned}
t_e^{(1)} &= \max(\sqrt{x_2}M_B, 1/b_1, 1/b_2)\ , \ \ t_e^{(2)} = \max(\sqrt{x_1}M_B, 1/b_1, 1/b_2)\ , \\
t_a^{(1)} &= \max(\sqrt{x_3}M_B, 1/b_2, 1/b_3)\ , \ \ t_a^{(2)} = \max(\sqrt{1-x_2}M_B, 1/b_2, 1/b_3)\ .
\end{aligned} \tag{19}$$

The Wilson coefficients are given by

$$\begin{aligned}
a_e(t) &= \left(C_3 + \frac{C_4}{N_c}\right) + \left(C_4 + \frac{C_3}{N_c}\right) + \left(C_5 + \frac{C_6}{N_c}\right) \\
&\quad -\frac{1}{2}\left(C_7 + \frac{C_8}{N_c}\right) - \frac{1}{2}\left(C_9 + \frac{C_{10}}{N_c}\right) - \frac{1}{2}\left(C_{10} + \frac{C_9}{N_c}\right)\ , \\
a_{a4}(t) &= \left(C_4 + \frac{C_3}{N_c}\right) + \frac{3}{2}e_q\left(C_{10} + \frac{C_9}{N_c}\right)\ , \\
a_{a6}(t) &= \left(C_6 + \frac{C_5}{N_c}\right) + \frac{3}{2}e_q\left(C_8 + \frac{C_7}{N_c}\right)\ , \\
a_2^T(t) &= C_2 + \frac{C_1}{N_c}\ ,
\end{aligned} \tag{20}$$

where a_2^T is the Wilson coefficient for the amplitude F_a^T. The hard functions, which are the Fourier transformation of the virtual quark propagator and the hard gluon propagator, are given by

$$\begin{aligned}
h_e(x_1, x_2, b_1, b_2) &= K_0\left(\sqrt{x_1 x_2}M_B b_1\right)\left[\theta(b_1 - b_2)K_0\left(\sqrt{x_2}M_B b_1\right)I_0\left(\sqrt{x_2}M_B b_2\right)\right. \\
&\quad \left. +\theta(b_2 - b_1)K_0\left(\sqrt{x_2}M_B b_2\right)I_0\left(\sqrt{x_2}M_B b_1\right)\right]\ ,
\end{aligned} \tag{21}$$

$$\begin{aligned}
h_a(x_2, x_3, b_2, b_3) &= \left(\frac{i\pi}{2}\right)^2 H_0^{(1)}\left(\sqrt{x_2 x_3}M_B b_2\right)\left[\theta(b_2 - b_3)H_0^{(1)}\left(\sqrt{x_3}M_B b_2\right)J_0\left(\sqrt{x_3}M_B b_3\right)\right. \\
&\quad \left. +\theta(b_3 - b_2)H_0^{(1)}\left(\sqrt{x_3}M_B b_3\right)J_0\left(\sqrt{x_3}M_B b_2\right)\right]\ .
\end{aligned} \tag{22}$$

B. Nonfactorizable Amplitudes



(a) (b) (c) (d)

FIG. 5. Feynman diagrams contributing to nonfactorizable amplitudes

The nonfactorizable amplitudes which come from Fig. 5 are written as

$$\mathcal{M}_e^P = \mathcal{M}_{e4}^P + \mathcal{M}_{e5}^P + \mathcal{M}_{e6}^P\ , \tag{23}$$

$$\mathcal{M}_{e4}^P = 16\pi C_F \frac{\sqrt{2N_c}}{N_c} M_B^4 \int_0^1 [dx] \int_0^\infty b_1 db_1 b_3 db_3 \phi_B(x_1, b_1)\phi_\phi(x_3)$$

$$
\begin{aligned}
&\times\Big\{ \left[(1-x_1-x_3)\phi_K^A(x_2) - r_K x_2\left(\phi_K^P(x_2)-\phi_K^T(x_2)\right)\right] E'_{e4}(t_d^{(1)})h_d^{(1)}(x_1,x_2,x_3,b_1,b_1,b_3) \\
&\quad + \left[(x_1-x_2-x_3)\phi_K^A(x_2) + r_K x_2\left(\phi_K^P(x_2)+\phi_K^T(x_2)\right)\right] E'_{e4}(t_d^{(2)})h_d^{(2)}(x_1,x_2,x_3,b_1,b_1,b_3)\Big\} \ ,
\end{aligned} \tag{24}
$$

$$
\begin{aligned}
\mathcal{M}_{e5}^P &= 16\pi C_F \frac{\sqrt{2N_c}}{N_c} M_B^4 \int_0^1 [dx] \int_0^\infty b_1 db_1 b_3 db_3 \phi_B(x_1,b_1)\phi_\phi(x_3) \\
&\times\Big\{ \left[(1-x_1+x_2-x_3)\phi_K^A(x_2) - r_K x_2\left(\phi_K^P(x_2)+\phi_K^T(x_2)\right)\right] E'_{e5}(t_d^{(1)})h_d^{(1)}(x_1,x_2,x_3,b_1,b_1,b_3) \\
&\quad + \left[(x_1-x_3)\phi_K^A(x_2) + r_K x_2\left(\phi_K^P(x_2)-\phi_K^T(x_2)\right)\right] E'_{e5}(t_d^{(2)})h_d^{(2)}(x_1,x_2,x_3,b_1,b_1,b_3)\Big\} \ ,
\end{aligned} \tag{25}
$$

$$
\begin{aligned}
\mathcal{M}_{e6}^P &= 16\pi C_F \frac{\sqrt{2N_c}}{N_c} M_B^4 r_\phi \int_0^1 [dx] \int_0^\infty b_1 db_1 b_3 db_3 \phi_B(x_1,b_1) \\
&\times\Big\{ \left[(1-x_1-x_3)\phi_K^A(x_2)\left(\phi_\phi^t(x_3)+\phi_\phi^s(x_3)\right)\right. \\
&\quad +r_K(1-x_1-x_3)\left(\phi_K^P(x_2)-\phi_K^T(x_2)\right)\left(\phi_\phi^t(x_3)+\phi_\phi^s(x_3)\right) \\
&\quad \left. -r_K x_2\left(\phi_K^P(x_2)+\phi_K^T(x_2)\right)\left(\phi_\phi^t(x_3)-\phi_\phi^s(x_3)\right)\right] E'_{e6}(t_d^{(1)})h_d^{(1)}(x_1,x_2,x_3,b_1,b_1,b_3) \\
&- \left[(x_1-x_3)\phi_K^A(x_2)\left(\phi_\phi^t(x_3)-\phi_\phi^s(x_3)\right) + r_K(x_1-x_3)\left(\phi_K^P(x_2)-\phi_K^T(x_2)\right)\left(\phi_\phi^t(x_3)-\phi_\phi^s(x_3)\right)\right. \\
&\quad \left. +r_K x_2\left(\phi_K^P(x_2)+\phi_K^T(x_2)\right)\left(\phi_\phi^t(x_3)+\phi_\phi^s(x_3)\right)\right] E'_{e6}(t_d^{(2)})h_d^{(2)}(x_1,x_2,x_3,b_1,b_1,b_3)\Big\} \ .
\end{aligned} \tag{26}
$$

$\mathcal{M}_{e3}^P$ is added to $\mathcal{M}_{e4}^P$ term, because $\mathcal{M}_{e3}^P$ is the same as $\mathcal{M}_{e4}^P$ without the Wilson coefficient.

$$
\mathcal{M}_a^P = \mathcal{M}_{a4}^P + \mathcal{M}_{a6}^P \ , \tag{27}
$$

$$
\begin{aligned}
\mathcal{M}_{a4}^P &= 16\pi C_F \frac{\sqrt{2N_c}}{N_c} M_B^4 \int_0^1 [dx] \int_0^\infty b_1 db_1 b_3 db_3 \phi_B(x_1,b_1) \\
&\times\Big\{ \left[(1-x_2)\phi_K^A(x_2)\phi_\phi(x_3) + r_K r_\phi\left((1-x_2)\left(\phi_K^P(x_2)-\phi_K^T(x_2)\right)\left(\phi_\phi^t(x_3)+\phi_\phi^s(x_3)\right)\right.\right. \\
&\quad \left.\left. +(x_1-x_3)\left(\phi_K^P(x_2)+\phi_K^T(x_2)\right)\left(\phi_\phi^t(x_3)-\phi_\phi^s(x_3)\right)\right)\right] E'_{a4}(t_f^{(1)})h_f^{(1)}(x_1,x_2,x_3,b_1,b_3,b_3) \\
&- \left[(x_1+x_3)\phi_K^A(x_2)\phi_\phi(x_3)\right. \\
&\quad +r_K r_\phi\left(4\phi_K^P(x_2)\phi_\phi^s(x_3) - (1-x_1-x_3)\left(\phi_K^P(x_2)-\phi_K^T(x_2)\right)\left(\phi_\phi^t(x_3)+\phi_\phi^s(x_3)\right)\right. \\
&\quad \left.\left. +x_2\left(\phi_K^P(x_2)+\phi_K^T(x_2)\right)\left(\phi_\phi^t(x_3)-\phi_\phi^s(x_3)\right)\right)\right] E'_{a4}(t_f^{(2)})h_f^{(2)}(x_1,x_2,x_3,b_1,b_3,b_3)\Big\} \ ,
\end{aligned} \tag{28}
$$

$$
\begin{aligned}
\mathcal{M}_{a6}^P &= 16\pi C_F \frac{\sqrt{2N_c}}{N_c} M_B^4 \int_0^1 [dx] \int_0^\infty b_1 db_1 b_3 db_3 \phi_B(x_1,b_1) \\
&\times\Big\{ \Big[- r_K(1-x_2)\left(\phi_K^P(x_2)-\phi_K^T(x_2)\right)\phi_\phi(x_3) \\
&\quad -r_\phi(x_1-x_3)\phi_K^A(x_2)\left(\phi_\phi^t(x_3)+\phi_\phi^s(x_3)\right)\Big] E'_{a6}(t_f^{(1)})h_f^{(1)}(x_1,x_2,x_3,b_1,b_3,b_3) \\
&+\Big[- r_K(1+x_2)\left(\phi_K^P(x_2)-\phi_K^T(x_2)\right)\phi_\phi(x_3) \\
&\quad -r_\phi(-2+x_1+x_3)\phi_K^A(x_2)\left(\phi_\phi^t(x_3)+\phi_\phi^s(x_3)\right)\Big] E'_{a6}(t_f^{(2)})h_f^{(2)}(x_1,x_2,x_3,b_1,b_3,b_3)\Big\} \ .
\end{aligned} \tag{29}
$$

The tree amplitude $\mathcal{M}_a^T$ is the same as $\mathcal{M}_{a4}$, but the Wilson coefficient is different. The evolution factors are defined by

$$
E'_{ei}(t) = \alpha_s(t) a'_{ei}(t) \exp[-S(t)|_{b_2=b_1}] \ , \quad E'_{ai}(t) = \alpha_s(t) a'_{ai}(t) \exp[-S(t)|_{b_2=b_3}] \ . \tag{30}
$$

The hard scales t are defined by

$$t_d^{(i)} = \max(DM_B, \sqrt{|D_i^2|}M_B, 1/b_1, 1/b_3)\ ,\ \ t_f^{(i)} = \max(FM_B, \sqrt{|F_i^2|}M_B, 1/b_1, 1/b_3)\ . \tag{31}$$

The Wilson coefficients are given by

$$\begin{aligned} a'_{e4}(t) &= C_3 + C_4 - \frac{1}{2}\left(C_9 + C_{10}\right)\ , & a'_{e5}(t) &= C_6 - \frac{1}{2}C_8\ , \\ a'_{e6}(t) &= C_5 - \frac{1}{2}C_7\ , & a'_{a4}(t) &= C_3 + \frac{3}{2}e_q C_9\ , \\ a'_{a6}(t) &= C_5 + \frac{3}{2}e_q C_7\ , & a^T(t) &= C_1. \end{aligned} \tag{32}$$

The hard functions are given by

$$\begin{aligned} h_d^{(j)} &= \left[\theta(b_1-b_3)K_0\left(DM_Bb_1\right)I_0\left(DM_Bb_3\right) + \theta(b_3-b_1)K_0\left(DM_Bb_3\right)I_0\left(DM_Bb_1\right)\right] \\ &\qquad \times K_0(D_jM_Bb_3) \qquad\qquad \text{for } D_j^2 \geq 0 \\ &\qquad \times \frac{i\pi}{2}H_0^{(1)}(\sqrt{|D_j^2|}M_Bb_3) \quad \text{for } D_j^2 \leq 0\ , \end{aligned} \tag{33}$$

$$\begin{aligned} h_f^{(j)} &= \frac{i\pi}{2}\left[\theta(b_1-b_3)H_0^{(1)}\left(FM_Bb_1\right)J_0\left(FM_Bb_3\right) + \theta(b_3-b_1)H_0^{(1)}\left(FM_Bb_3\right)J_0\left(FM_Bb_1\right)\right] \\ &\qquad \times K_0(F_jM_Bb_1) \qquad\qquad \text{for } F_j^2 \geq 0 \\ &\qquad \times \frac{i\pi}{2}H_0^{(1)}(\sqrt{|F_j^2|}M_Bb_1) \quad \text{for } F_j^2 \leq 0\ , \end{aligned} \tag{34}$$

with the variables

$$\begin{aligned} D^2 &= x_1x_2\ , & F^2 &= (1-x_2)x_3\ , \\ D_1^2 &= \{x_1-(1-x_3)\}x_2\ , & F_1^2 &= (x_1-x_3)(1-x_2)\ , \\ D_2^2 &= (x_1-x_3)x_2\ , & F_2^2 &= x_1+x_3+(1-x_1-x_3)(1-x_2)\ . \end{aligned} \tag{35}$$

IV. NUMERICAL RESULTS

We summarize the parameters which is used in the numerical analysis of our calculation.

We use the model of the B meson wave function written as

$$\phi_B(x,b) = N_Bx^2(1-x)^2\exp\left[-\frac{1}{2}\left(\frac{xM_B}{\omega_B}\right)^2 - \frac{\omega_B^2b^2}{2}\right]\ , \tag{36}$$

where $\omega_B = 0.40$ GeV [22]. N_B is determined from normalization condition Eq. (B5).

The K meson wave functions are given as

$$\phi_K^A(x) = \frac{f_K}{2\sqrt{2N_c}}6x(1-x)\left[1 + 3a_1(1-2x) + \frac{3}{2}a_2\{5(1-2x)^2-1\}\right]\ , \tag{37}$$

$$\begin{aligned} \phi_K^P(x) = \frac{f_K}{2\sqrt{2N_c}}\Bigg[&1 + \frac{1}{2}\left(30\eta_3 - \frac{5}{2}\rho_K^2\right)\{3(1-2x)^2-1\} \\ &-\frac{1}{8}\left(3\eta_3\omega_3 + \frac{27}{20}\rho_K^2 + \frac{81}{10}\rho_K^2a_2\right)\{3 - 30(1-2x)^2 + 35(1-2x)^4\}\Bigg]\ , \end{aligned} \tag{38}$$

$$\phi_K^T(x) = \frac{f_K}{2\sqrt{2N_c}}(1-2x)\left[1 + 6\left(5\eta_3 - \frac{1}{2}\eta_3\omega_3 - \frac{7}{20}\rho_K^2 - \frac{3}{5}\rho_K^2a_2\right)(1-10x+10x^2)\right]\ . \tag{39}$$

where $\rho_K = (M_d + m_s)/M_K$ [23] [24]. The parameters of these wave functions are given as $a_1 = 0.17,\ a_2 = 0.20,\ \eta_3 = 0.015,\ \omega_3 = -3.0$ where renormalization scale is 1 GeV. We fix the parameters as above values.

The ϕ meson wave functions are given as

$$\phi_\phi(x) = \frac{f_\phi}{2\sqrt{2N_c}} 6x(1-x) , \tag{40}$$

$$\begin{aligned}\phi^t_\phi(x) = \frac{f^T_\phi}{2\sqrt{2N_c}} \Bigg[& 3(1-2x)^2 + \frac{35}{4}\zeta^T_3 \{3 - 30(1-2x)^2 + 35(1-2x)^4\} \\ & + \frac{3}{2}\delta_+ \left\{ 1 - (1-2x)\log\frac{1-x}{x} \right\} \Bigg] ,\end{aligned} \tag{41}$$

$$\phi^s_\phi(x) = \frac{f^T_\phi}{4\sqrt{2N_c}} \left[(1-2x)\left(6 + 9\delta_+ + 140\zeta^T_3 - 1400\zeta^T_3 x + 1400\zeta^T_3 x^2\right) + 3\delta_+ \log\frac{x}{1-x} \right] , \tag{42}$$

where $\zeta^T_3 = 0.024,\ \delta_+ = 0.46$ [25]. We fix the parameters as above values, since the numerical results is insensitive to these parameters.

We use the Wolfenstein parameters for the CKM matrix elements $A = 0.819,\ \lambda = 0.2196,\ R_b \equiv \sqrt{\rho^2 + \eta^2} = 0.38$ [26], and choose the angle $\phi_3 = \pi/2$ [6]. In addition, we use the following parameters.

$$\begin{gathered} M_B = 5.28 \text{ GeV}, \quad M_K = 0.49 \text{ GeV}, \quad M_\phi = 1.02 \text{ GeV}, \quad m_{0K} = 1.70 \text{ GeV}, \\ f_B = 190 \text{ MeV}, \quad f_K = 160 \text{ MeV}, \quad f_\phi = 237 \text{ MeV}, \quad f^T_\phi = 215 \text{ MeV}, \\ \tau_{B^0} = 1.55 \times 10^{-12} \text{ sec}, \quad \tau_{B^\pm} = 1.65 \times 10^{-12} \text{ sec} , \quad \Lambda^{(4)}_{\rm QCD} = 0.250 \text{ GeV}. \end{gathered}$$

We show the numerical results of each amplitude for the $B^0 \to \phi K^0$ and $B^\pm \to \phi K^\pm$ decays in Table I. The factorizable penguin amplitude F^P_e is dominant contribution to the $B \to \phi K$ decays. The factorizable annihilation penguin amplitude F^P_a generates large strong phase. In the $B^\pm \to \phi K^\pm$ modes, there are the contribution from $f_B F^T_a$ and M^T_a. These tree amplitudes contribute a few percent to the whole amplitude, since the CKM matrix elements related to the tree amplitudes are very small.

We predict the branching ratios for the $B \to \phi K$ decays as

$$\text{Br}(B^0 \to \phi K^0) = \left| \frac{f_B f_K}{190 \text{ MeV } 160 \text{ MeV}} \right|^2 \times \left(9.43 \times 10^{-6}\right) , \tag{43}$$

$$\text{Br}(B^\pm \to \phi K^\pm) = \left| \frac{f_B f_K}{190 \text{ MeV } 160 \text{ MeV}} \right|^2 \times \left(10.1 \times 10^{-6}\right) . \tag{44}$$

The current experimental values are summarized in Table II. The values which are predicted in PQCD are consistent with the current experimental data. However, our branching ratios have the theoretical error from the $O(\alpha_s^2)$ corrections, the higher twist corrections, and the error of input parameters. Large uncertainties come from the meson decay constants, the shape parameter ω_B, and m_{0K}. These parameters are fixed from the other modes($B \to K\pi,\ D\pi\ \pi\pi$, etc.). We try to vary ω_B from 0.36 GeV to 0.44 GeV, then we obtain $\text{Br}(B^\pm \to \phi K^\pm) = (7.54 \sim 13.9) \times 10^{-6}$. Next, we set $\omega_B = 0.40$ and try to vary m_{0K} from 1.40 GeV to 1.80 GeV, then we obtain $\text{Br}(B^\pm \to \phi K^\pm) = (6.65 \sim 11.4) \times 10^{-6}$.

In the naive factorization assumption, the branching ratio is very sensitive to the effective number of colors $N_c^{\rm eff}$. If we set $N_c^{\rm eff}$=3, then the branching ratio is about 4.5×10^{-6} where the scale of the Wilson coefficient is taken to $M_B/2$ and F^{BK} is 0.38 from the BSW model. Our predicted values are larger than those obtains from the naive factorization assumption. This is due to the enhancement of the Wilson coefficient. In PQCD approach, the scale of the Wilson coefficients, which is equal to the hard scale t, can reach lower values than $M_B/2$ in case of the naive factorization assumption, and the Wilson coefficients are then enhanced [6]. Therefore, our branching ratios are enhanced over those obtained from the naive factorization assumption.

In the QCD-improved factorization, the branching ratios for the $B \to \phi K$ decays are predicted as $\mathrm{Br}(B^0 \to \phi K^0) = (4.0^{+2.9}_{-1.4}) \times 10^{-6}$ and $\mathrm{Br}(B^- \to \phi K^-) = (4.3^{+3.0}_{-1.4}) \times 10^{-6}$ with annihilation effect [12]. Our predicted values are larger than these values.

We also compute the direct CP asymmetry defined as

$$A_{CP}^{\mathrm{dir}} = \frac{|A^+|^2 - |A^-|^2}{|A^+|^2 + |A^-|^2} \ . \tag{45}$$

If the amplitude of the $B^\pm \to \phi K^\pm$ decay can be written as

$$A^+ = V_{ts} V_{tb}^* e^{i\delta_P} A_P - V_{us} V_{ub}^* e^{i\delta_T} A_T \ , \tag{46}$$

$$A^- = V_{ts}^* V_{tb} e^{i\delta_P} A_P - V_{us}^* V_{ub} e^{i\delta_T} A_T \ , \tag{47}$$

where the indices T and P denote the tree and penguin respectively, δ_T and δ_P are the strong phase, and the amplitude A_T and A_P are real, then the direct CP asymmetry is given by

$$A_{CP}^{\mathrm{dir}} = \frac{2\lambda^2 R_b \sin\phi_3 \sin(\delta_P - \delta_T) \frac{A_T}{A_P}}{1 + \lambda^4 R_b \left(\frac{A_T}{A_P}\right)^2} \ . \tag{48}$$

Our results of the direct CP asymmetry is about few percent. The reason why the direct CP asymmetry is very small is that the CKM matrix element for the tree amplitude is much smaller than for the penguin amplitude.

V. SUMMARY

In this paper, we calculate the $B^0 \to \phi K^0$ and $B^\pm \to \phi K^\pm$ decays in PQCD approach. Our predicted branching ratios agree with the current experimental data and are larger than the values obtained by the naive factorization assumption and the QCD-improved factorization. This is because the Wilson coefficients related to penguin operators are enhanced dynamically in PQCD.

In the PQCD approach, we can calculate the strong phase from nonfactorizable amplitudes and annihilation amplitudes explicitly. We can then calculate the direct CP asymmetry of this modes, however this asymmetry is very small since the CKM matrix element for the tree amplitude is much smaller than for the penguin amplitude.

Note added:

After this work has been completed, we become aware of a similar calculation by Chen, Keum and Li [27]. Our results are in agreement.

Acknowledgment

The topic of this research was suggested by Professor A.I. Sanda. The author thanks our PQCD group members: Y.Y. Keum, E. Kou, T. Kurimoto, H-n. Li, T. Morozumi, R. Sinha, K. Ukai for useful discussions. The author thanks JSPS for partial support. The work was supported in part by Grant-in Aid for Special Project Research (Physics of CP violation); Grant-in Aid for Scientific Research from the Ministry of Education, Science and Culture of Japan.

APPENDIX A: WILSON COEFFICIENTS

The Wilson coefficients for the $m_b = 4.8$ GeV are

$$\begin{aligned} &C_1(m_b) = -2.693\times 10^{-1}\ , &&C_2(m_b) = 1.123\ , \\ &C_3(m_b) = 1.243\times 10^{-2}\ , &&C_4(m_b) = -2.665\times 10^{-2}\ , \\ &C_5(m_b) = 7.753\times 10^{-3}\ , &&C_6(m_b) = -3.256\times 10^{-2}\ , \\ &C_7(m_b) = 3.426\times 10^{-4}\ , &&C_8(m_b) = 3.131\times 10^{-4}\ , \\ &C_9(m_b) = -9.732\times 10^{-3}\ , &&C_{10}(m_b) = 2.217\times 10^{-3}\ . \end{aligned} \tag{A1}$$

APPENDIX B: B MESON WAVE FUNCTIONS

We consider the B meson wave function $\Phi^{(in)}_{B,\alpha\beta,ij} = \langle 0|\bar{b}_{\beta j}(0)d_{\alpha i}(z)|B(P_1)\rangle$ where the indices α and β are the spin indices, i and j are the color indices. This wave function can be decomposed into some components with different spin structure. The B meson wave function is then decomposed to be

$$\Phi^{(in)}_{B,\alpha\beta,ij} = \frac{i\delta_{ij}}{\sqrt{2N_c}}\int dx_1 d^2\mathbf{k}_{1T} e^{-i(x_1P_1^- z^+ - \mathbf{k}_{1T}\mathbf{z}_T)}\left[(\not{P}_1\gamma_5)\phi_B^A(x_1,\mathbf{k}_{1T}) + \gamma_5\phi_B^P(x_1,\mathbf{k}_{1T})\right]_{\alpha\beta}\ , \tag{B1}$$

where N_c is color's degree of freedom. The axial-vector wave function $\phi_B^A(x_1,\mathbf{k}_{1T})$ is defined by

$$P_{1\mu}\int dx_1 d^2\mathbf{k}_{1T} e^{-i(x_1P_1^- z^+ - \mathbf{k}_{1T}\mathbf{z}_T)}\phi_B^A(x_1,\mathbf{k}_{1T}) = \frac{-i}{2\sqrt{2N_c}}\langle 0|\bar{b}(0)\gamma_5\gamma_\mu d(z)|\bar{b}(P_1)\rangle\ , \tag{B2}$$

and the pseudo-scalar wave function $\phi_B^P(x_1,\mathbf{k}_{1T})$ is defined by

$$\int dx_1 d^2\mathbf{k}_{1T} e^{-i(x_1P_1^- z^+ - \mathbf{k}_{1T}\mathbf{z}_T)}\phi_B^P(x_1,\mathbf{k}_{1T}) = \frac{-i}{2\sqrt{2N_c}}\langle 0|\bar{b}(0)\gamma_5 b(z)|\bar{d}(P_1)\rangle\ . \tag{B3}$$

Using HQET, we can denote $\phi_B^P(x_1,\mathbf{k}_{1T}) = M_B\phi_B^A(x_1,\mathbf{k}_{1T})$. The B meson wave function is then written by

$$\Phi^{(in)}_{B,\alpha\beta,ij} = \frac{i\delta_{ij}}{\sqrt{2N_c}}\int dx_1 d^2\mathbf{k}_{1T} e^{-i(x_1P_1^- z^+ - \mathbf{k}_{1T}\mathbf{z}_T)}\left[(\not{P}_1 + M_B)\gamma_5\phi_B(x_1,\mathbf{k}_{1T})\right]_{\alpha\beta}\ . \tag{B4}$$

The normalization of this wave function is given by

$$\int_0^1 dx_1\phi_B(x_1, b_1 = 0) = \frac{f_B}{2\sqrt{2N_c}}\ , \tag{B5}$$

where b_1 is the conjugate space of k_1, and f_B is the decay constant of the B meson.

APPENDIX C: K MESON WAVE FUNCTIONS

The K meson wave function is defined that $\Phi^{(in)}_{K,\alpha\beta,ij} = \langle 0|\bar{s}_{\beta j}(0)d_{\alpha i}(z)|K(P_2)\rangle$, $z = (0, z^-, \mathbf{0}_T)$ is the coordinate of the d quark. $\Phi^{(in)}_{K,\alpha\beta,ij}$ can be decomposed into some components with different spin structure $\gamma_5\gamma_\mu$, γ_5, $\gamma_5\sigma_{\mu\nu}$, $\cdots$. These matrix elements are calculated using the twist expansion in the light-cone QCD sum rules. The twist expansion is the expansion in terms of m_{0K} where $m_{0K} = M_K^2/(m_d + m_s)$. The leading twist terms are independent on m_{0K}, the twist-3 and twist-4 terms are proportional to m_{0K} and m_{0K}^2 respectively. Up to twist-3, the matrix elements are given as [23] [24]

$$\langle 0|\bar{s}(0)\gamma_5\gamma_\mu d(z)|K(P_2)\rangle = -if_K P_{2\mu}\int_0^1 dx_2 e^{-ix_2P_2\cdot z}\phi_v(x_2)\ , \tag{C1}$$

$$\langle 0|\bar{s}(0)\gamma_5 d(z)|K(P_2)\rangle = -if_K m_{0K}\int_0^1 dx_2 e^{-ix_2P_2\cdot z}\phi_p(x_2)\ , \tag{C2}$$

$$\begin{aligned}\langle 0|\bar{s}(0)\gamma_5\sigma_{\mu\nu} d(z)|K(P_2)\rangle &= \frac{i}{6}f_K m_{0K}\left(1-\frac{M_K^2}{m_{0K}^2}\right)(P_{2\mu}z_\nu - P_{2\nu}z_\mu)\int_0^1 dx_2 e^{-ix_2P_2\cdot z}\phi_\sigma(x_2)\\ &= \frac{1}{6}f_K m_{0K}(v_\mu n_\nu - v_\nu n_\mu)\int_0^1 dx_2 e^{-ix_2P_2\cdot z}\frac{d}{dx_2}\phi_\sigma(x_2)\ ,\end{aligned} \tag{C3}$$

where $v^\mu = P_2^\mu/P_2^+$ and $n^\mu = z^\mu/z^- = (0,1,\mathbf{0}_T)$. $\phi_v(x_2)$ is the leading twist wave function, $\phi_p(x_2)$ and $d/dx_2\phi_\sigma(x_2)$ are the twist-3 wave functions. These wave functions are normalized as

$$\int_0^1 dx_2\phi_v(x_2) = 1\ ,\ \int_0^1 dx_2\phi_p(x_2) = 1\ ,\ \int_0^1 dx_2\frac{d}{dx_2}\phi_\sigma(x_2) = 0\ . \tag{C4}$$

We define the following new wave functions,

$$\phi_K^A(x_2) = \frac{f_K}{2\sqrt{2N_c}}\phi_v(x_2)\ ,\ \phi_K^P(x_2) = \frac{f_K}{2\sqrt{2N_c}}\phi_p(x_2)\ ,\ \phi_K^T(x_2) = \frac{f_K}{12\sqrt{2N_c}}\frac{d}{dx_2}\phi_\sigma(x_2)\ . \tag{C5}$$

The K meson wave functions are then written as

$$\Phi^{(in)}_{K,\alpha\beta,ij} = \frac{-i\delta_{ij}}{\sqrt{2N_c}}\int_0^1 dx_2 e^{-ix_2P_2\cdot z}\left[\not{P}_2\gamma_5\phi_K^A(x_2) + m_{0K}\gamma_5\phi_K^P(x_2) + m_{0K}(\not{n}\,\not{v}-1)\gamma_5\phi_K^T(x_2)\right]_{\alpha\beta}\ . \tag{C6}$$

APPENDIX D: ϕ MESON WAVE FUNCTIONS

The ϕ meson wave function has much terms than the B meson and K meson wave functions, since the ϕ meson is a vector meson. The wave function $\Phi^{(in)}_{\phi,\alpha\beta,ij} = \langle 0|\bar{s}_{\beta j}(0)s_{\alpha i}(z)|\phi(P_3)\rangle$ can be decomposed into some components with different spin structure γ_μ, $\gamma_\mu\gamma_5$, $\sigma_{\mu\nu}$, I, $\cdots$. The components are calculated using the twist expansion in the light-cone QCD sum rules. In this case, the twist expansion is the expansion in terms of M_ϕ. Up to twist-3, the components are given as [25]

$$\begin{aligned}\langle 0|\bar{s}(0)\gamma_\mu s(z)|\phi(P_3)\rangle &= f_\phi M_\phi\int_0^1 dx_3 e^{-ix_3P_3\cdot z}\left[P_{3\mu}\frac{\epsilon_\phi\cdot z}{P_3\cdot z}\phi_{//}(x_3) + \epsilon^T_{\phi\mu}g_\perp^{(v)}(x_3)\right]\\ &= f_\phi M_\phi\int_0^1 dx_3 e^{-ix_3P_3\cdot z}\left[\epsilon_{\phi\mu}\phi_{//}(x_3) + \epsilon^T_{\phi\mu}g_\perp^{(v)}(x_3)\right]\ ,\end{aligned} \tag{D1}$$

$$\begin{aligned}\langle 0|\bar{s}(0)\gamma_\mu\gamma_5 s(z)|\phi(P_3)\rangle &= -\frac{1}{4}\left(f_\phi - f_\phi^T\frac{2m_s}{M_\phi}\right)M_\phi\epsilon_\mu{}^{\nu\alpha\beta}\epsilon^T_{\phi\nu}P_{3\alpha}z_\beta\int_0^1 dx_3 e^{-ix_3P_3\cdot z}g_\perp^{(a)}(x_3)\\ &= -\frac{1}{4}f_\phi M_\phi\epsilon_\mu{}^{\nu\alpha\beta}\epsilon^T_{\phi\nu}P_{3\alpha}z_\beta\int_0^1 dx_3 e^{-ix_3P_3\cdot z}g_\perp^{(a)}(x_3)\ ,\end{aligned} \tag{D2}$$

$$\begin{aligned}\langle 0|\bar{s}(0)\sigma_{\mu\nu} s(z)|\phi(P_3)\rangle &= if_\phi^T\int_0^1 dx_3 e^{-ix_3P_3\cdot z}\Big[\left(\epsilon^T_{\phi\mu}P_{3\nu} - \epsilon^T_{\phi\nu}P_{3\mu}\right)\phi_\perp(x_3)\\ &\qquad\qquad +(P_{3\mu}z_\nu - P_{3\nu}z_\mu)\frac{\epsilon_\phi\cdot z}{(P_3\cdot z)^2}M_\phi^2 h_{//}^{(t)}(x_3)\Big]\\ &= if_\phi^T\int_0^1 dx_3 e^{-ix_3P_3\cdot z}\Big[\left(\epsilon^T_{\phi\mu}P_{3\nu} - \epsilon^T_{\phi\nu}P_{3\mu}\right)\phi_\perp(x_3)\\ &\qquad\qquad +(\epsilon_{\phi\mu}P_{3\nu} - \epsilon_{\phi\nu}P_{3\mu})h_{//}^{(t)}(x_3)\Big]\ ,\end{aligned} \tag{D3}$$

$$\begin{aligned}\langle 0|\bar{s}(0) I s(z)|\phi(P_3)\rangle &= \frac{i}{2}\left(f_\phi^T - f_\phi \frac{2m_s}{M_\phi}\right)\epsilon_\phi \cdot z M_\phi^2 \int_0^1 dx_3 e^{-ix_3 P_3\cdot z} h_{//}^{(s)}(x_3) \\ &= -f_\phi^T M_\phi \int_0^1 dx_3 e^{-ix_3 P_3\cdot z}\frac{1}{2}\frac{d}{dx_3} h_{//}^{(s)}(x_3)\ , \end{aligned} \tag{D4}$$

where $n_\mu = z_\mu/z^+$,$v_\mu = P_{3\mu}/P_3^-$. $\phi_{//}(x_3)$ and $\phi_\perp(x_3)$ are the leading twist wave functions, $h_{//}^{(t)}(x_3)$, $\frac{d}{dx_3}h_{//}^{(s)}(x_3)$, $g_\perp^{(v)}(x_3)$ and $g_\perp^{(a)}(x_3)$ are the twist-3 wave functions. These wave functions satisfy the normalization conditions

$$\begin{aligned} &\int_0^1 dx_3 \phi_{//}(x_3) = 1\ ,\ \int_0^1 dx_3 h_{//}^{(t)}(x_3) = 1\ ,\ \int_0^1 dx_3 \frac{d}{dx_3} h_{//}^{(s)}(x_3) = 0\ , \\ &\int_0^1 dx_3 \phi_\perp(x_3) = 1\ ,\ \int_0^1 dx_3 g_\perp^{(v)}(x_3) = 1\ ,\quad \int_0^1 dx_3 g_\perp^{(a)}(x_3) = 1\ . \end{aligned} \tag{D5}$$

We can neglect the wave functions which are proportional to the transverse polarization vector ϵ_ϕ^T, because these terms do not appear in our calculations. We define the following new wave functions,

$$\phi_\phi(x_3) = \frac{f_\phi}{2\sqrt{2N_c}}\phi_{//}(x_3)\ ,\ \phi_\phi^t(x_3) = \frac{f_\phi^T}{2\sqrt{2N_c}} h_{//}^{(t)}(x_3)\ ,\ \phi_\phi^s(x_3) = \frac{f_\phi^T}{4\sqrt{2N_c}}\frac{d}{dx_3} h_{//}^{(s)}(x_3)\ . \tag{D6}$$

The ϕ meson wave functions are then written as

$$\Phi_{\phi,\alpha\beta,ij}^{(in)} = \frac{\delta_{ij}}{\sqrt{2N_c}}\int_0^1 dx_3 e^{-ix_3 P_3\cdot z}\left[M_\phi \not\epsilon_\phi \phi_\phi(x_3) + \not P_3 \not\epsilon_\phi \phi_\phi^t(x_3) + M_\phi \phi_\phi^s(x_3)\right]_{\alpha\beta}\ . \tag{D7}$$

It must be noted that this expression is dependent on how to define the sign of ϵ_ϕ.

We considered the wave functions for an initial state. In this study, we need the K meson and ϕ meson wave functions for a final state. The wave function for a final state is given as $\Phi^{(out)} = \gamma_0 \Phi^{(in)\dagger}\gamma_0$.

---

[1] G. Cavoto on behalf of the BaBar Collaboration, hep-ex/0105018.

[2] A. Bozek for the BELLE Collaboration, talk presented at the International Europhysics Conference on High Energy Physics, Budapest, HUNGARY, July 12-18, 2001.

[3] CLEO Collaboration, Phys. Rev. Lett. **86**, 3718 (2001).

[4] For example, see I.I. Bigi and A.I. Sanda, *CP Violation* (2000, Cambridge).

[5] M. Bauer, B. Stech, M. Wirbel, Z. Phys. C **34**, 103 (1987); Z. Phys. C **29**, 637 (1985).

[6] Y.Y. Keum, H-n. Li and A.I. Sanda, Phys. Lett. B **504**, 6(2001); Phys. Rev. D **63**, 054008 (2001).

[7] C.D. Lü, K. Ukai, and M.Z. Yang, Phys. Rev. D **63**, 074009 (2001).

[8] C.D. Lü and M.Z. Yang, hep-ph/0011238.

[9] C.H. Chang and H-n. Li, Phys. Rev. D **63**, 014003 (2001).

[10] E. Kou and A.I. Sanda, hep-ph/0106159.

[11] X.G. He, J.P. Ma and C.Y. Wu, Phys. Rev. D **63**, 094004 (2001).

[12] H.Y. Cheng and K.C. Yang, hep-ph/0012152.

[13] M. Kobayashi and T. Maskawa, Prog. Th. Phys. **49**, 652 (1973).

[14] G. Buchalla, A.J. Buras and M.E. Lautenbacher, Review of Modern Physics, **68**, 1125 (1996).

[15] C.H. Chang and H-n. Li, Phys. Rev. D **55**, 5577 (1997).

[16] T.W. Yeh and H-n. Li, Phys. Rev. D **56**, 1615 (1997).

[17] J.C. Collins and D.E. Soper, Nucl. Phys. B**193**, 381 (1981).

[18] J. Botts and G. Sterman, Nucl. Phys. **B325**, 62 (1989).

[19] H-n. Li, Phys. Rev. D **52**, 3958 (1995).

[20] H-n. Li, hep-ph/0102013.

[21] T. Kurimoto, H-n. Li and A.I. Sanda, hep-ph/0105003.

[22] M. Bauer and M. Wirbel, Z. Phys. C**42**, 671 (1989).

[23] P. Ball, JHEP 09, 005 (1998).

[24] P. Ball, JHEP 01, 010 (1999).

[25] P. Ball, V.M. Braun, Y. Koike, and K. Tanaka, Nucl. Phys. B**529**, 323 (1998).

[26] Particle Data Group, Eur. Phys. J. C**15** 1 (2000).

[27] C.H. Chen, Y.Y. Keum and H-n. Li, hep-ph/0107165.

	$B^0 \to \phi K^0$	$B^\pm \to \phi K^\pm$
$f_\phi F_e^P$	-1.03×10^{-1}	-1.03×10^{-1}
$f_B F_a^P$	$6.45 \times 10^{-3} + \ i\ 4.28 \times 10^{-2}$	$6.17 \times 10^{-3} + \ i\ 4.20 \times 10^{-2}$
M_e^P	$5.24 \times 10^{-3} - \ i\ 3.61 \times 10^{-3}$	$5.24 \times 10^{-3} - \ i\ 3.61 \times 10^{-3}$
M_a^P	$-8.03 \times 10^{-4} - \ i\ 1.73 \times 10^{-3}$	$-6.56 \times 10^{-4} - \ i\ 7.22 \times 10^{-4}$
$f_B F_a^T$		$-1.11 \times 10^{-1} - \ i\ 3.75 \times 10^{-2}$
M_a^T		$1.60 \times 10^{-2} + \ i\ 2.77 \times 10^{-2}$

TABLE I. Contribution to the $B^0 \to \phi K^0$ and $B^\pm \to \phi K^\pm$ decays from each amplitude.

	$\mathrm{Br}(B^0 \to \phi K^0)$	$\mathrm{Br}(B^\pm \to \phi K^\pm)$
BaBar	$(8.1^{+3.1}_{-2.5} \pm 0.8) \times 10^{-6}$	$(7.7^{+1.6}_{-1.4} \pm 0.8) \times 10^{-6}$
BELLE	$(8.7^{+3.8}_{-3.0} \pm 1.5) \times 10^{-6}$	$(10.6^{+2.1}_{-1.9} \pm 2.2) \times 10^{-6}$
CLEO	$< 12.3 \times 10^{-6}$	$(5.5^{+2.1}_{-1.8} \pm 0.6) \times 10^{-6}$

TABLE II. The experimental data of the $B \to \phi K$ branching ratios from the BaBar [1], the BELLE [2] and the CLEO [3]